

<b>DISEASE</b>	<b>PATIENT SHALL BE EXCLUDED FROM SCHOOL</b>	<b>CONTACT</b>
PEDICULOSIS (Head Lice)	Until appropriate treatment has commenced. Treatment should be repeated after 7 days.	Not to be excluded. Close contacts should be inspected regularly for infestation.
PERTUSIS (Whooping Cough)	For 4 weeks, or until a medical certificate of recovery is produced.	Not to be excluded.
RINGWORM	Until appropriate treatment has commenced, supported when requested by a medical certificate.	Not to be excluded.
RUBELLA (German Measles)	Until fully recovered, and at least 4 days from the onset of the rash.	Not to be excluded.
SCABIES	Until appropriate treatment has commenced, supported when requested by a medical certificate.	Not to be excluded.
STREPTOCOCCAL INFECTIONS (including Scarlet Fever)	Until a medical certificate of recovery is produced.	Not to be excluded.7680.
CHICKENPOX	Until fully recovered. Note: some remaining scabs are not an indication for continued exclusion.	Not to be excluded.
VIRAL HEPATITIS	Until a medical certificate of recovery is produced, or on subsidence of symptoms, but not before 7 days after onset of jaundice.	Not to be excluded.
IMPETIGO (School Sores)	Until sores have healed. The child may be allowed to return, provided that appropriate treatment is being applied and that sores on exposed surfaces, such as scalp, face, hands and legs are properly covered with occlusive dressings.	Not to be excluded.
MEASLES	For at least 7 days from the appearance of the rash, or a medical certificate of recovery is produced.	To be excluded for 13 days after last contact, unless medical documentation of prior infection or immunisation is provided.
MENINGOCOCCAL INFECTION	Until a medical certificate of recovery is produced.	If domiciliary, to be excluded until they have been receiving appropriate chemotherapy for at least 48 hours.
MUMPS	Until fully recovered.	Not to be excluded.